arXiv:1303.5865v1 [math.MG] 23 Mar 2013

Geometric construction of adding and dissection of triangles.

E. Mieczkowski*

Abstract

This article is a continuation and development of the 1st chapter of the article [6]. The set of segments, each of the next is n times bigger than the first one is a simple geometric interpretation of the set $\mathbb{N}$ of natural numbers. In this paper we investigate the opposite situation. We give algebraic structure to the set of similar triangles with parallel sides. We present geometric construction of adding of triangles and use it to dissection of triangles into 15 triangles of different sides.

Introduction

We can see at the set of integer numbers $\mathbb{Z}$ as the one-dimensional module over the set $\mathbb{Z}$. Geometric interpretation of any number $n \in \mathbb{Z}$ is vector n times bigger than the vector represented by number 1 or -1. Let us take the fixed triangle and the set of similar triangles with the sides parallel and n times bigger than the fixed one's sides. We try to give geometric structure to this set. In the second section we give a geometric construction of adding of triangles. In the third section we use this construction to write dissection of the triangle for 15 triangles of different sides.

1 Basic concepts

Let us set the triangle on the plan $\mathbb{R}^2$ and let us denote it by the symbol $\langle 1 \rangle$. Then the similar triangle with sides parallel to the sides of $\langle 1 \rangle$ and n times

---

*Department of Mathematics, Gdynia Maritime University, Gdynia, Poland; E-mail: `edek@am.gdynia.pl`.

bigger sides (we denote this triangle by the symbol $\langle n\rangle$) is built from $\frac{n(n+1)}{2}$ triangles $\langle 1\rangle$ and $\frac{n(n-1)}{2}$ symmetrical to triangle $\langle 1\rangle$ in relation to any side, triangle denoted by symbol $\langle -1\rangle$ (see Fig. 1). So



Fig. 1.

$$\langle n\rangle = \frac{n(n+1)}{2}\langle 1\rangle + \frac{n(n-1)}{2}\langle -1\rangle \tag{1}$$

Let us mark by the symbol $\langle -n\rangle$ the triangle similar to the triangle $\langle -1\rangle$ with parallel and n times bigger sides. Then

$$\begin{aligned}\langle -n\rangle &= \frac{n(n-1)}{2}\langle 1\rangle + \frac{n(n+1)}{2}\langle -1\rangle\\ &= \frac{(-n)(-n+1)}{2}\langle 1\rangle + \frac{(-n)(-n-1)}{2}\langle -1\rangle\end{aligned}$$

Each point of the plane $\mathbb{R}^2$ we will denote by $\langle 0\rangle$. Later each triangle $\langle \pm n\rangle$, where $n \in \mathbb{N}$ we will denote by the symbol $\langle n\rangle$, where $n \in \mathbb{Z}$.
The symbol $-\langle n\rangle$ denotes the triangle, which lying on the triangle $\langle n\rangle$ gives the empty set.
The set $\mathbb{N}_2 = \{\pm\langle n\rangle; n \in \mathbb{Z}\}$ is the subset of the ring

$$\mathbb{P}_2(\mathbb{Z}) = \{(x, y) = x\langle 1\rangle + y\langle -1\rangle, x, y \in \mathbb{Z}\}$$

with addition

$$\forall x_1, x_2, y_1, y_2 \in \mathbb{Z} \quad (x_1, y_1) + (x_2, y_2) = (x_1 + x_2, y_1 + y_2)$$

and commutative multiplication

$$(x_1, y_1) \cdot (x_2, y_2) = (x_1 x_2 + y_1 y_2, x_1 y_2 + x_2 y_1)$$

and the neutral element $(0, 0) = \langle 0 \rangle$ of addition and neutral element $(1, 0) = \langle 1 \rangle$ of multiplication.
It is easy to see that

$$\langle -1 \rangle^2 = \langle 1 \rangle$$

$$\langle 1 \rangle \cdot \langle -1 \rangle = \langle -1 \rangle \cdot \langle 1 \rangle = \langle -1 \rangle$$

and $\forall n, m \in \mathbb{Z}$

$$\langle n \rangle \cdot \langle m \rangle = \langle nm \rangle$$

Let us transform

$$\langle n \rangle = \frac{n^2 + n}{2} \langle 1 \rangle + \frac{n^2 - n}{2} \langle -1 \rangle = n^2 \frac{\langle 1 \rangle + \langle -1 \rangle}{2} + n \frac{\langle 1 \rangle - \langle -1 \rangle}{2}$$

It is easy to show that elements $\frac{\langle 1 \rangle + \langle -1 \rangle}{2} = A_2$ and $\frac{\langle 1 \rangle - \langle -1 \rangle}{2} = A_1$ are orthogonal. Let us denote $A_2 = \binom{1}{0}$, $\quad A_1 = \binom{0}{1}$. Then

$$\langle n \rangle = n^2 A_2 + n A_1 \tag{2}$$

we can call the orthogonal form of the triangle $\langle n \rangle$. Using the linear combination of the elements of the set $\mathbb{N}_2$ we can write different geometric figures (Fig. 2, 3).
From the description of Fig. 2, 3 we can see that the coefficient of A_2 means the number of all triangles $\langle \pm 1 \rangle$ in the given figure. The second coefficient of A_1 equals to the difference in the lengths of the parallel sides (the unit side length equals to the length of the parallel side of the triangle $\langle 1 \rangle$). So if the triangle $\langle 1 \rangle$ is equilateral or not, the length of each side is equal to 1.

Let us put the addition closed in $\mathbb{N}_2$:

$\forall n, k, l \in \mathbb{Z}$

$$\langle n + k + l \rangle = \langle n + k \rangle + \langle n + l \rangle + \langle k + l \rangle - \langle n \rangle - \langle k \rangle - \langle l \rangle \tag{3}$$



$\langle n\rangle - \langle k\rangle =$
$(n^2-k^2)A_2 + (n-k)A_1,$
$n > k > 0$

$\langle -n\rangle - \langle -k\rangle =$
$(n^2-k^2)A_2 - (n-k)A_1,$
$n > k > 0$

Fig. 2.



$\langle n+k\rangle - \langle n\rangle - \langle k\rangle =$
$= 2nkA_2,$
$n > k > 0$

$\langle n+k+l+t\rangle - \langle n\rangle - \langle k\rangle - \langle l\rangle =$
$= xA_2 + tA_1,$
$n, k, l, t \in \mathbb{N}$

Fig. 3.

The operation is well-defined. This results from orthogonal form of $\langle n \rangle$ and the truth of the below condition:
$\forall n, k, l \in \mathbb{Z} \quad \forall i = 1, 2$

$$(n+k+l)^i = (n+k)^i + (n+l)^i + (k+l)^i - (n)^i - (k)^i - (l)^i.$$

The operation (3) has the simple geometric interpretation for $n, k, l > 0$ (Fig. 4).



$\triangle ADG = \langle n+k+l \rangle$

$\triangle BDF = \langle n+k \rangle$ $\quad \triangle ACH = \langle n+l \rangle$ $\quad \triangle EGI = \langle k+l \rangle$

$\triangle BCJ = \langle n \rangle$ $\quad \triangle EFJ = \langle k \rangle$ $\quad \triangle HIJ = \langle l \rangle$

Fig. 4.

Let us consider the equation (3) for the concrete numbers

$$\langle 4 \rangle = \langle 1+1+2 \rangle = 2\langle 3 \rangle + \langle 2 \rangle - \langle 2 \rangle - 2\langle 1 \rangle \tag{4}$$

After reduction we get

$$\langle 4 \rangle = \langle 1+1+2 \rangle = 2\langle 3 \rangle - 2\langle 1 \rangle \tag{5}$$

From the arithmetic point of view the equation (5) is true. But it easy to see that we can not build the triangle $\langle 4 \rangle$ using only two triangles $\langle 3 \rangle$ and two triangles $-\langle 1 \rangle$. We need the triangles $\langle 2 \rangle$ and $-\langle 2 \rangle$ too. They are not reducible to the empty set because they do not lie on one another. (Fig. 5).



Fig. 5.

Definition 1. *The equation* $\langle n\rangle = \sum_j \alpha_j \langle n_j\rangle$*, where* $\alpha_j \in \{-1, 1\}$ *is true in the geometric sense if we can build the triangle* $\langle n\rangle$ *from the elements* $\alpha_j\langle n_j\rangle$*.*

Definition 2. *The equation* $\langle n\rangle = \sum_j \alpha_j \langle n_j\rangle$*, where* $\alpha_j \in \{-1, 1\}$ *is true in the arithmetic sense if the equations* $n^i = \sum_j \alpha_j n_j^i$*, where* $\alpha_j \in \{-1, 1\}$*,* $i = 1, 2$ *hold.*

Corollary 1. *If the equation is true in the geometric sense, it is also true in the arithmetic sense.*

So we can say that the equation (4) is true in the geometric sense, while the equation (5) only in the arithmetic sense.

2 Generalization of the equation (3)

We can generalize the equation (3). Namely

$$\begin{aligned} &\forall n, k, l, t \in \mathbb{Z} \\ &\qquad \langle n+k+l+t\rangle = \langle n+k+t\rangle + \langle n+l+t\rangle + \langle k+l+t\rangle \\ &\qquad\qquad -\langle n+t\rangle - \langle k+t\rangle - \langle l+t\rangle + \langle t\rangle \end{aligned} \tag{6}$$

The equation (6) is true in arithmetic sense because the below equation holds for $i = 2, 1$.

$$\begin{aligned} (n+k+l+t)^i &= (n+k+t)^i + (n+l+t)^i + (k+l+t)^i \\ &\quad -(n+t)^i - (k+t)^i - (l+t)^i + t^i. \end{aligned}$$

The Fig. 6 shows that for $n, k, l, t > 0$ the equation (6) is true in the geometric sense as well.



Fig. 6. Interpretation of the equation (6).

In order to prove that the equation (6) holds for any n, k, l, t in the geometric sense we will create a certain geometric construction of this operation. Let us fix the ordered successive components n, k, l of the sum $\langle n + k + l + t\rangle$ extend or shorten the triangle $\langle t\rangle$ in the directions I, II, III, or I ', II ', III', depending on whether the numbers n, k, l are positive or negative (Fig. 7). Therefore the equation (6) should be properly written as



Fig. 7.

$$
\begin{aligned}
&\forall n,k,l,t \in \mathbb{Z} \\
&\langle (n+k+l+t) \rangle = \langle n+k+0+t \rangle + \langle n+0+l+t \rangle + \langle 0+k+l+t \rangle \\
&\qquad - \langle n+0+0+t \rangle - \langle 0+k+0+t \rangle - \langle 0+0+l+t \rangle \\
&\qquad + \langle 0+0+0+t \rangle
\end{aligned}
\tag{7}
$$

Below are further examples of the creation of new triangles from the triangle $\langle t \rangle$ (Fig. 8-11). Triangle $\langle t \rangle$ has vertices ABC, and the new one has vertices $A'B'C'$.



$\langle n+0+0+t \rangle$
$n, t > 0$

$\langle n+0+0+t \rangle$
$0 < n+t < t$

Fig. 8.



$\langle n+0+0+t \rangle$
$n < -t < 0$

$\langle 0+k+0+t \rangle$
$0 < k+t < t$

Fig. 9.

Since the first component will be generally denoted by the letter n, k second, third l, so if it will not lead to confusion, we will write $\langle n+t \rangle$, $\langle k+t \rangle$, $\langle l+t \rangle$ instead of respectively $\langle n+0+0+t \rangle$, $\langle 0+k+0+t \rangle$, $\langle 0+0+l+t \rangle$. Similarly we will write $\langle n+k+t \rangle$, $\langle n+k+t \rangle$, $\langle k+l+t \rangle$ instead of $\langle n+k+0+t \rangle$, $\langle n+k+0+t \rangle$, $\langle 0+k+l+t \rangle$.



$\langle n + k + 0 + t\rangle$

$n > 0,\ 0 < k + t < 0$

$\langle n + k + 0 + t\rangle$

$0 < -t < n,\ k < 0$

Fig. 10.



$\langle n + k + l + t\rangle$

$n > 0,\ 0 < k + t < 0,\ l < -t < 0$

Fig. 11.

Triangles on Fig. 11 have the following descriptions.
$\triangle AB_nC_n = \langle n+t\rangle$, $\triangle BC_kA_k = \langle k+t\rangle$, $\triangle CA_lB_l = \langle l+t\rangle$,
$\triangle A_kB_nC' = \langle n+k+t\rangle$, $\triangle A'B_lC_k = \langle k+l+t\rangle$, $\triangle A_lB'C_n = \langle n+l+t\rangle$.

It's easy to follow, that they satisfy with the triangles ABC and $A'B'C'$ equation (6) in geometric sense. It should also be noted that the expression $\langle n+k+l+t\rangle$ is the whole figure of Fig. 11, constructed of eight triangles, while $\langle (n{+}k{+}l{+}t)\rangle = \langle 0{+}0{+}0{+}(n{+}k{+}l{+}t)\rangle$ means only the triangle $A'B'C'$.

Theorem 1. *$\forall n,k,l,t \in \mathbb{Z}$ the equation (7) is true in the geometric sense.*

Proof. Let us note the equations (8)-(10) have the same geometric interpretation as the equation (7).

$$\begin{aligned}
\langle (n+k+t)\rangle &= \langle n+k+(-l)+(l+t)\rangle \\
&= \langle n+k+0+(l+t)\rangle + \langle n+0+(-l)+(l+t)\rangle \\
&\quad +\langle 0+k+(-l)+(l+t)\rangle \\
&\quad -\langle n+0+0+(l+t)\rangle - \langle 0+k+0+(l+t)\rangle \\
&\quad -\langle 0+0+(-l)+(l+t)\rangle + \langle (l+t)\rangle \\
&= \langle n+k+l+t\rangle + \langle n+t\rangle + \langle k+t\rangle \\
&\quad -\langle n+l+t\rangle - \langle k+l+t\rangle - \langle t\rangle + \langle l+t\rangle
\end{aligned} \tag{8}$$

$$\begin{aligned}
\langle (n+t)\rangle &= \langle n+(-k)+(-l)+(k+l+t)\rangle \\
&= \langle n+(-k)+0+(k+l+t)\rangle + \langle n+0+(-l)+(k+l+t)\rangle \\
&\quad +\langle 0+(-k)+(-l)+(k+l+t)\rangle \\
&\quad -\langle n+0+0+(k+l+t)\rangle - \langle 0+(-k)+0+(k+l+t)\rangle \\
&\quad -\langle 0+0+(-l)+(l+k+t)\rangle + \langle 0+0+0+(k+l+t)\rangle \\
&= \langle n+l+t\rangle + \langle n+k+t\rangle + \langle t\rangle \\
&\quad -\langle n+k+l+t\rangle - \langle l+t\rangle - \langle k+t\rangle + \langle k+l+t\rangle
\end{aligned} \tag{9}$$

$$\begin{aligned}
\langle t\rangle &= \langle(-n)+(-k)+(-l)+(n+k+l+t)\rangle \\
&= \langle 0+(-k)+(-l)+(n+k+l+t)\rangle \\
&\quad +\langle(-n)+0+(-l)+(n+k+l+t)\rangle \\
&\quad +\langle(-n)+(-k)+0+(n+k+l+t)\rangle \\
&\quad -\langle 0+0+(-l)+(n+k+l+t)\rangle \\
&\quad -\langle 0+(-k)+0+(n+k+l+t)\rangle \\
&\quad -\langle(-n)+0+0+(n+k+l+t)\rangle+\langle 0+0+0+(n+k+l+t)\rangle \\
&= \langle n+t\rangle+\langle k+t\rangle+\langle l+t\rangle \\
&\quad -\langle n+k+t\rangle-\langle n+l+t\rangle-\langle k+l+t\rangle+\langle n+k+l+t\rangle \qquad (10)
\end{aligned}$$

So if in the equation (7) one element l is negative we can replace the equation (7) by equation (8). If in the equation (7) two elements k, l are negative we can replace the equation (7) by equation (9). If the equation (7) has three elements n, k, l negative we take the equation (10). So it is sufficient to consider the cases $n > 0, k > 0, l > 0$ and any t. We have 10 following cases.

(1) $t > 0$, (Fig. 6)
In next cases $t < 0$.

(2) $n+t > 0, k+t > 0, l+t > 0$, (Fig. 12).

(3) $n+t > 0, k+t > 0, l+t < 0$, (Fig. 13).

(4) $n+t > 0, k+t < 0, l+t < 0, k+l+t > 0$, (Fig. 14).

(5) $n+t > 0, k+l+t < 0$, (Fig. 15).

(6) $n+t < 0, k+t < 0, l+t < 0, n+k+t > 0, k+l+t > 0, n+l+t > 0$, (Fig. 16).

(7) $n+t < 0, n+k+t > 0, k+l+t < 0, n+l+t > 0$, (Fig. 17).

(8) $n+k+t > 0, k+l+t < 0, n+l+t < 0$, (Fig. 18).

(9) $n+k+t < 0, k+l+t < 0, n+l+t < 0, n+k+l+t > 0$, (Fig. 19).

(10) $n+k+l+t < 0$, (Fig. 20).

The proof is based on reviewing each figure and founding that by using the components of the equation (7), we always get the triangle $\langle (n+k+l+t) \rangle$ from the triangle $\langle t \rangle$. On Fig. 12-20 the triangle $\langle (n+k+l+t) \rangle$ is denoted by $\langle s \rangle$.



Fig. 12.



Fig. 13.



Fig. 14.



Fig. 15.



Fig. 16.



Fig. 17.



Fig. 18.



Fig. 19.



Fig. 20.

□

The equations (8)-(10) are equivalent to the subtraction of the additive group G. That means, if a, b, c are elements of the group G and satisfy the equation $a = b + c$ then there exists an element $(-c) \in G$ fulfilling $b = a + (-c)$.
It is easy to see that we can not get similar equations from the formula (3).

Theorem 2. $\forall n, k, l, t \in \mathbb{Z}$
If $n+k+l+t = 0$, then the triangles $\langle t\rangle$ and $\langle n+k+l+t\rangle$ from the equation (7) are congruent.

Proof. Since $n + k + l + t = 0$, so $\langle n + k + l + t\rangle = \langle t\rangle$. This completes the proof. □

Corollary 2. $\forall n, k, l \in \mathbb{Z}$
If $n + k + l = 0$, then the equation $\langle n + k + l + 0\rangle = \langle 0\rangle$ is a translation of the point $\langle 0\rangle$ in another point.

Theorem 3. *If $n \neq n' \vee k \neq k' \vee l \neq l'$,
then $\langle n + k + l + t\rangle \neq \langle (n' + k' + l' + t)\rangle$ in the geometric sense.*

Proof. The proof follows from the geometric construction of the operation (7). ☐

The set $\mathbb{N}_2$ can be extended to the set

$$\mathbb{Q}_2 = \{\pm\langle q\rangle = \pm(q^2A_2 + qA_1);\ q \in \mathbb{Q}\} \subset P_2(\mathbb{Q}),$$

where $\mathbb{Q}$ is a set of rational numbers and $P_2(\mathbb{Q})$ is the ring with the same operations as in the ring $P_2(\mathbb{Z})$. Of course, the operation (7) is also true in the set $\mathbb{Q}_2$ in arithmetic and geometric sense.

3 Dissection of triangles into triangles

We will use the set $\mathbb{N}_2$ to write solution of the following problem. It is known [1, 2] that a square can be dissected into at least 21 squares of different sides. It is also known [3]-[5] that a triangle can be dissected into at least 15 similar triangles of different sides. At the same, it is considered that the triangles, one of which is a mirror image of the other are different. On (Fig. 21) we have one of two possible optimum dissection of the triangle.



Fig. 21.

This dissection can be written by using 7 times the equation (7).

$$\begin{aligned}
\langle 39\rangle &= \langle 19+12+20-12\rangle \\
&= \langle 19+12+0-12\rangle + \langle 0+12+20-12\rangle + \langle 19+0+20-12\rangle \\
&\quad -\langle 19+0+0-12\rangle - \langle 0+12+0-12\rangle - \langle 0+0+20-12\rangle \\
&\quad +\langle 0+0+0-12\rangle \\
&= \langle 19\rangle + \langle 20\rangle + \langle 27\rangle - \cancel{\langle 7\rangle}_1 - \cancel{\langle 8\rangle}_2 + \langle -12\rangle, \qquad (11)
\end{aligned}$$

where

$$\langle 27\rangle = \langle 11+16+11-11\rangle = \langle 16_1\rangle + \langle 11\rangle + \langle 16_2\rangle - \bcancel{\langle 5\rangle}_3 + \langle -11\rangle, \qquad (12)$$

where

$$\langle 16_1\rangle = \langle 7+7+9-7\rangle = \bcancel{\langle 7\rangle}_1 + \langle 9_1\rangle + \langle 9_2\rangle - \bcancel{\langle 2\rangle}_4 + \langle -7\rangle, \qquad (13)$$

where

$$\langle 9_2\rangle = \langle 2+7+2-2\rangle = \langle 7_1\rangle + \bcancel{\langle 2\rangle}_4 + \langle 7_2\rangle - \bcancel{\langle 5\rangle}_5 + \langle -2\rangle, \qquad (14)$$

where

$$\langle 7_2\rangle = \langle 5+5+2-5\rangle = \bcancel{\langle 5\rangle}_5 + \langle 2_1\rangle + \bcancel{\langle 2_2\rangle}_7 - \bcancel{\langle -3\rangle}_6 + \langle -5\rangle, \qquad (15)$$

where

$$\langle 16_2\rangle = \langle 8+8+8-8\rangle = \langle 8_1\rangle + \langle 8_2\rangle + \bcancel{\langle 8_3\rangle}_2 + \langle -8\rangle, \qquad (16)$$

where

$$\langle 8_1\rangle = \langle 3+5+3-3\rangle = \bcancel{\langle 5_1\rangle}_3 + \langle 5_2\rangle + \langle 3\rangle - \bcancel{\langle 2\rangle}_7 + \bcancel{\langle -3\rangle}_6. \qquad (17)$$

It should be noted that the triangles in order to be reduced have to be of different signs and lie one on the other. We can see the equations (11)-(17) on Fig. 22.



Fig. 22.

The equations (11)-(17) are not the only way to write this dissection of the triangle. The equations (18)-(24) and (Fig. 23) gives another way to write.

$$\begin{aligned}
\langle 39\rangle &= \langle 19+19+20-19\rangle \\
&= \langle 19+19+0-19\rangle + \langle 0+19+20-19\rangle + \langle 19+0+20-19\rangle \\
&\quad -\langle 19+0+0-19\rangle - \langle 0+19+0-19\rangle - \langle 0+0+20-19\rangle \\
&\quad +\langle 0+0+0-19\rangle \\
&= \langle 19_1\rangle + \langle 20_1\rangle + \bcancel{\langle 20_2\rangle}_6 - \bcancel{\langle 1\rangle}_1 + \langle -19\rangle, \qquad (18)
\end{aligned}$$

where

$$\langle -19\rangle = \langle -12-7-7+7\rangle = \langle 12_1\rangle + \langle -7\rangle + \langle 12_2\rangle - \bcancel{\langle -5\rangle}_2 + \langle 7\rangle, \qquad (19)$$

where

$$\langle -12_1\rangle = \langle -5-7-5+5\rangle = \bcancel{\langle -7_1\rangle}_7 + \langle -7_2\rangle + \bcancel{\langle -5\rangle}_2 - \bcancel{\langle -2\rangle}_3 + \langle 5\rangle, \qquad (20)$$

where

$$\langle -7_2\rangle = \langle -5-2-2+2\rangle = \langle -5_1\rangle + \langle -2\rangle + \langle -5_2\rangle - \bcancel{\langle -3\rangle}_4 + \langle 2\rangle, \qquad (21)$$

where

$$\langle -5_1\rangle = \langle -3-2-3+3\rangle = \bcancel{\langle -2_1\rangle}_3 + \langle -2_2\rangle + \bcancel{\langle -3\rangle}_4 - \bcancel{\langle 1\rangle}_5 + \langle 3\rangle, \qquad (22)$$

where

$$\langle -2_2\rangle = \langle -11+9-11+11\rangle = \langle 9_1\rangle + \langle 9_2\rangle + \langle -11\rangle - \bcancel{\langle 20\rangle}_6 + \langle 11\rangle, \qquad (23)$$

where

$$\langle 9_2\rangle = \langle 8+1+8-8\rangle = \bcancel{\langle 1_1\rangle}_5 + \bcancel{\langle 1_2\rangle}_1 + \langle 8\rangle - \bcancel{\langle -7\rangle}_7 + \langle -8\rangle, \qquad (24)$$


⟨39⟩ = ⟨19 + 19 + 20 − 19⟩
⟨19⟩
⟨20⟩
⟨−19⟩
⟨−12⟩
−⟨−5⟩
⟨7⟩
⟨−7⟩
⟨1⟩
⟨20⟩
⟨−12⟩
⟨−7⟩ = ⟨−5 − 2 − 2 + 2⟩
⟨−5⟩
⟨−5⟩
−⟨−3⟩
⟨2⟩
⟨−2⟩
⟨−5⟩
⟨5⟩
⟨−7⟩
⟨−19⟩ = ⟨−12 − 7 − 7 + 7⟩
⟨−3⟩
⟨−7⟩
−⟨−2⟩
⟨3⟩
⟨−2⟩
⟨−2⟩
⟨9⟩
⟨−12⟩ = ⟨−5 − 7 − 5 + 5⟩
⟨−5⟩ = ⟨−3 − 2 − 3 + 3⟩
−⟨1⟩
⟨−2⟩
⟨−2⟩ = ⟨−11 + 9 − 11 + 11⟩
⟨−11⟩
⟨9⟩
−⟨−7⟩
⟨11⟩
⟨1⟩
⟨−8⟩
⟨1⟩
⟨8⟩
⟨9⟩ = ⟨8 + 1 + 8 − 8⟩

Fig. 23.

References

[1] A.J.W Duijvestijn, *Simple perfect squared square of lowest order* J. Combin. Theory, Ser. B, 25 (1978), pp. 240243.

[2] I. Gambini, *A method for cutting squares into distinct squares*, Discrete Applied Mathematics, Volume 98, Issues 12 (1999), pp. 6580.

[3] W. T. Tutte, *Dissections into Equilateral Triangles*, The Mathematical Gardner, edited by David Klarner. Published by Van Nostrand Reinhold (1981).

[4] W. T. Tutte, *The Dissection of Equilateral Triangles into Equilateral Triangles*, Proc. Cambridge Phil. Soc 44 (1948), pp. 463-482.

[5] A. Drápal, C. Hämäläinen published *An enumeration of equilateral triangle dissections*, Discrete Applied Mathematics, Volume 158, Issue 14 (2010).

[6] E. Mieczkowski, *The arithmetic of solids*, Preprint arXiv:1204.2219, 6 Apr 2012v1.